

GONALITY AND CLIFFORD INDEX OF CURVES ON $K3$ SURFACES

ANDREAS LEOPOLD KNUTSEN

ABSTRACT. We show that every possible value for the Clifford index and gonality of a curve of a given genus on a $K3$ surface occurs.

1. INTRODUCTION

Curves on $K3$ surfaces have been objects of intense study throughout the years. In particular, in connection with the famous conjecture of Green [Gr], Green and Lazarsfeld proved in [G-L] that the Clifford index is the same for all smooth curves in a complete linear system on a $K3$ surface. The original conjecture by Harris and Mumford was that the gonality should be the same for all smooth curves in a linear system, but this was proved to be false by a counterexample of Donagi and Morrison [D-M]. If the line bundle is ample, then this is indeed the only counterexample, by a result of Ciliberto and Pareschi [C-P].

In [E-L-M-S] curves on $K3$ surfaces are used to construct exceptional curves of any Clifford dimension. In our recent paper [Kn1] we relate the Clifford index and gonality of curves to the higher order embeddings of the surface.

The aim of this note is to show that for $g \geq 3$, genus g curves of any possible Clifford index and gonality can be found on some $K3$ surface. More precisely, we will show the following statements:

Theorem 1.1. (a) *Let g and c be integers such that $g \geq 4$ and $0 \leq c \leq \lfloor \frac{g-1}{2} \rfloor$. Then there exists a $K3$ surface containing a smooth curve of genus g and Clifford index c .*

(b) *Let g and k be integers such that $g \geq 3$ and $2 \leq k \leq \lfloor \frac{g+3}{2} \rfloor$. Then there exists a $K3$ surface containing a smooth curve of genus g and gonality k .*

Recall that a smooth curve C of genus $g \geq 2$ is said to have *gonality* k if C possesses a g_k^1 but no g_{k-1}^1 .

If A is a line bundle on C , then the *Clifford index* of A is the integer

$$\text{Cliff } A = \deg A - 2(h^0(A) - 1).$$

When $g \geq 4$, the *Clifford index* of C itself is defined as

$$\text{Cliff } C = \min\{\text{Cliff } A \mid h^0(A) \geq 2, h^1(A) \geq 2\}.$$

Note that we always have $\text{Cliff } C \leq \text{gon } C - 2$. Furthermore, we get from Brill-Noether theory that the gonality of C satisfies $\text{gon } C \leq \lfloor \frac{g+3}{2} \rfloor$, whence $\text{Cliff } C \leq \lfloor \frac{g-1}{2} \rfloor$. For the general curve of genus g , we have $\text{gon } C = \lfloor \frac{g+3}{2} \rfloor$ and $\text{Cliff } C = \lfloor \frac{g-1}{2} \rfloor$.

We say that a line bundle A on C *contributes to the Clifford index of C* if $h^0(A) \geq 2$ and $h^1(A) \geq 2$ and that it *computes the Clifford index of C* if in addition $\text{Cliff } C = \text{Cliff } A$.

It is classically known that for a fixed genus, curves of any possible gonality occur. By Ballico's result [Ba], the same is true for the Clifford index. Our result then gives a new

1991 *Mathematics Subject Classification.* 14J28.

Key words and phrases. curves, $K3$ surfaces, Clifford index, gonality.

proof of these facts, with the additional information that one can always construct such curves on $K3$ surfaces.

The central tool in this paper is the already mentioned famous result of Green and Lazarsfeld [G-L], which states that if L is a base point free line bundle on a $K3$ surface S , then $\text{Cliff } C$ is constant for all smooth irreducible $C \in |L|$, and if $\text{Cliff } C < \lfloor \frac{g-1}{2} \rfloor$, then there exists a line bundle M on S such that $M_C := M \otimes \mathcal{O}_C$ computes the Clifford index of C for all smooth irreducible $C \in |L|$. (Note that since $(L - M) \otimes \mathcal{O}_C \simeq \omega_C \otimes M_C^{-1}$, the result is symmetric in M and $L - M$.)

Other tools are classical results on linear systems on $K3$ surfaces [SD] and lattice theory of $K3$ surfaces [Mo].

For other results on the Clifford index and gonality of smooth curves on $K3$ surfaces we refer to [C-P], [E-L-M-S] and [Kn1].

A *curve* is always reduced and irreducible.

The author would like to thank Gavril Farkas for interesting conversations.

Remark 1.2. We adopt the same convention as in [G-L], and set $\text{Cliff } C = 0$ for C of genus 2 or hyperelliptic of genus 3, and $\text{Cliff } C = 1$ for C non-hyperelliptic of genus 3. The result of Green and Lazarsfeld is then still valid.

2. PROOF OF THE MAIN THEOREM

The theorem is an immediate consequence of the following:

Proposition 2.1. *Let d and g be integers such that $g \geq 3$ and $2 \leq d \leq \lfloor \frac{g+3}{2} \rfloor$, and let S be a $K3$ surface with $\text{Pic } S = \mathbf{Z}L \oplus \mathbf{Z}E$, where $L^2 = 2(g-1)$, $E.L = d$ and $E^2 = 0$. Then L is base point free and for any smooth curve $C \in |L|$ we have*

$$\text{Cliff } C = d - 2 \leq \lfloor \frac{g-1}{2} \rfloor$$

and $\text{Cliff } C$ is computed by the pencil $\mathcal{O}_C(E)$.

To prove this proposition, we first need the following basic existence result:

Lemma 2.2. *Let $g \geq 3$ and $d \geq 2$ be integers. Then there exists a $K3$ surface S with $\text{Pic } S = \mathbf{Z}L \oplus \mathbf{Z}E$, such that L is base point free and E is a smooth curve, $L^2 = 2(g-1)$, $E.L = d$ and $E^2 = 0$.*

Proof. By [Kn2, Prop. 4.2], we can find a $K3$ surface S with $\text{Pic } S = \mathbf{Z}L \oplus \mathbf{Z}E$, with intersection matrix

$$\begin{bmatrix} L^2 & L.E \\ E.L & E^2 \end{bmatrix} = \begin{bmatrix} 2(g-1) & d \\ d & 0 \end{bmatrix}$$

and such that L is nef. This is a consequence of the lattice theory in [Mo], which again follows from the surjectivity of the period map. If L is not base point free, there exists by standard results on linear systems on $K3$ surfaces (see e.g. [SD, 2.7] or [Kn1, Thm. 1.1]) a curve B such that $B^2 = 0$ and $B.L = 1$. An easy calculation, writing $B \sim xL + yE$ for two integers x and y , shows that this is impossible. By [Kn2, Proposition 4.4], we have that $|E|$ contains a smooth curve. $\square$

Proof of Proposition 2.1. Let S , L and E be as in Lemma 2.2, with $d \leq \lfloor \frac{g+3}{2} \rfloor$. Note that since E is irreducible, we have $h^1(E) = 0$. Furthermore, Riemann-Roch gives $\chi(L - E) = \frac{1}{2}(L - E)^2 + 2 = g + 1 - d \geq \frac{g-1}{2} \geq 1$, whence by Serre duality either $h^0(L - E) \geq 1$ or $h^0(E - L) \geq 1$ (but not both simultaneously). Since $(E - L).L = d - 2g + 2 < 0$, we have $h^0(L - E) \geq 1$. If $g \geq 4$, we even have $h^0(L - E) \geq 2$.

By the cohomology of the short exact sequence

$$0 \longrightarrow \mathcal{O}_S(E - L) \longrightarrow \mathcal{O}_S(E) \longrightarrow \mathcal{O}_C(E) \longrightarrow 0,$$

where C is any smooth curve in $|L|$, we find that $h^0(\mathcal{O}_C(E)) \geq h^0(E) = 2$ and $h^1(\mathcal{O}_C(E)) = h^0(L - E)$, so $\mathcal{O}_C(E)$ contributes to the Clifford index of C (if $g \geq 4$) and

$$c := \text{Cliff } C \leq \text{Cliff } \mathcal{O}_C(E) \leq E.L - E^2 - 2 = d - 2 < \lfloor \frac{g-1}{2} \rfloor.$$

If $c = \lfloor \frac{g-1}{2} \rfloor$, there is nothing more to prove.

If $c < \lfloor \frac{g-1}{2} \rfloor$, then by [Ma, (2.3)] and [Kn1, Lemma 8.3] there exists an effective divisor D on S satisfying

- (1) $h^0(\mathcal{O}_C(D)) = h^0(D) \geq 2, \quad h^1(\mathcal{O}_C(D)) = h^0(L - D) \geq 2,$
- (2) $h^1(D) = h^1(L - D) = 0 \quad \text{and}$
- (3) $c = \text{Cliff } \mathcal{O}_C(D) = \text{Cliff } \mathcal{O}_C(L - D) = D.L - D^2 - 2$

(this is a consequence of the result of Green and Lazarsfeld [G-L] mentioned above).

Since both $D > 0$ and $L - D > 0$ and E is nef, we must have

$$D.E \geq 0 \text{ and } (L - D).E \geq 0.$$

Writing $D \sim xL + yE$ this is equivalent to

$$dx \geq 0 \text{ and } d(1 - x) \geq 0,$$

which gives $x = 0$ or 1 . These two cases give, respectively, $D = yE$ and $L - D = yE$. Since $h^1(yE) = y - 1$ by [SD, Prop. 2.6], it follows by (2) that $y = 1$ and $D \sim E$ or $L - D \sim E$. Hence $c = \text{Cliff } \mathcal{O}_C(E) = E.L - E^2 - 2 = d - 2$ by (3). By Riemann-Roch and (1) we also have $h^0(\mathcal{O}_C(E)) = h^0(E) = \frac{1}{2}E^2 + 2 = 2$, so $\mathcal{O}_C(E)$ is a pencil. $\square$

This concludes the proof of Theorem 1.1.

REFERENCES

- [Ba] E. Ballico, *On the Clifford index of algebraic curves*, Proc. Amer. Math. Soc. **97**, 217-218 (1986).
- [C-P] C. Ciliberto, G. Pareschi, *Pencils of minimal degree on curves on a $K3$ surface*, J. für die reine und angew. Math. **460** (1995), 15-36.
- [D-M] R. Donagi, D. R. Morrison, *Linear systems on $K3$ sections*, J. Diff. Geom. **29**, 49-64 (1989).
- [E-L-M-S] D. Eisenbud, H. Lange, G. Martens, F.-O. Schreyer, *The Clifford dimension of a projective curve*, Comp. Math. **72**, 173-204 (1989).
- [Gr] M. Green, *Koszul cohomology and the geometry of projective varieties*, J. Differ. Geom. **19**, 125-171 (1984).
- [G-L] M. Green, R. Lazarsfeld, *Special divisors on curves on a $K3$ surface*, Invent. Math. **89**, 357-370 (1987).
- [Kn1] A. L. Knutsen, *On k th order embeddings of $K3$ surfaces and Enriques surfaces*, Manuscr. Math. **104**, 211-237 (2001).
- [Kn2] A. L. Knutsen, *Smooth Curves on Projective $K3$ Surfaces*, to appear in Math. Scand., preprint available at math.AG/9805140.
- [Ma] G. Martens, *On curves on $K3$ surfaces*, in E. Ballico, C. Ciliberto (eds): *Algebraic curves and projective geometry*, Pproceedings, 1988, Lecture Notes in Math. 1389, Springer-Verlag (1989).
- [Mo] D. R. Morrison, *On $K3$ surfaces with large Picard number*, Invent. Math. **75**, 105-121 (1984).
- [SD] B. Saint-Donat, *Projective Models of $K3$ Surfaces*, Amer. J. Math. **96**, 602-639 (1974).